

### **CONSERVATION ENHANCEMENT ACTIVITY**

## E3281



# Forage harvest to reduce water quality impacts by utilization of excess soil nutrients

**Conservation Practice 328: Conservation Crop Rotation** 

**APPLICABLE LAND USE: Crop (Annual & Mixed); Crop (Perennial)** 

**RESOURCE CONCERN: Water** 

**ENHANCEMENT LIFE SPAN: 1 Year** 

#### **Enhancement Description**

Establish a forage crop (single species or mix) following a primary annual crop to take up excess soil nutrients. Select forage known to effectively utilize and scavenge nutrients. Forage shall be harvested for forage, but not be grazed or burned.

#### Criteria

- This enhancement is applicable on fields where excess soil nutrients cause or increase water quality degradation concerns. Presence of excess nutrients must be identified in recent soil tests or increased risk to water quality documented by risk assessment tool. (Refer to state specific guidance of options to maximize nutrient uptake in local climate and cropping systems)
- Forage species, seedbed preparation, seeding rates, seeding dates, seeding depths, fertility requirements, and planting methods will be consistent with applicable local criteria and soil/site conditions. (Refer to state specific lists of forage crops known to effectively utilize and scavenge nutrients)
- Select forage crop (single species or mix of two or more species) and planting dates
  which will not compete with the other crop(s) yield or harvest. If legumes are part
  of the forage mix, consider that this may add nutrients to the system.

E328I - Forage harvest to reduce water	August 2019	
quality impacts by utilization of excess soil	Montana Supplement March 2020	Page   1
Nutrients with Montana Supplement		



#### **United States Department of Agriculture**

- Select forage crop that is compatible with other components of the crop rotation and for its ability to efficiently scavenge and utilize excess soil nutrients, specifically nitrogen or phosphorous, whichever is identified as a potential risk to water quality. Nutrient uptake only occurs when a crop is actively growing. Therefore, it is imperative that the crops in rotation be planted as soon as possible after forage crop harvest (hay/balage/haylage/etc.) to maximize nutrient cycling and minimize offsite transport of nutrients.
- Determine method and timing of forage crop harvest to meet client objectives.
   Harvest the forage crop as late as practical to maximize plant biomass production and nutrient uptake.
- Ensure any herbicides used in the crop rotation are compatible with forage crop selections.
- Do not burn forage or residue.
- Do not graze forage crop.
- Reduce or maintain soil erosion from water and wind to below soil tolerance (T) level (average annual soil loss).



# **Documentation and Implementation Requirements**

# Participant will:

• Prior to implementation, provide NRCS with the current and planned crop rotation and field operation(s) used for each crop.

**Document excess nutrients identified in soil tests.** Soil tests should be taken as close to production crop harvest as possible.

Field	Soil Test Date	Nutrient (Nitrogen or Phosphorus)	Soil Test Nutrient R <mark>esu</mark> (ppm or lbs/ <mark>ac)</mark>

**Current Management Rotation** 

Field	Current Crops (in sequence)	Planting Date	Harvest I	Date
			V	

**Current Field Operations for Each Crop** 

Field	Сгор	Field Operation	Timing of Field Operation (month/year)
			The second second

E328I - Forage harvest to reduce water	August 2019	
quality impacts by utilization of excess soil	Montana Supplement March 2020	Page   3
Nutrients with Montana Supplement		

#### Planned Management Rotation including Forage Crop

Field	Planned Crops/Forage Crop (in sequence)	Planting Date	Harvest Date

#### **Planned Field Operations for Each Crop**

Field	Crop	Field Operation	Timing of Field Operation (month/year)

#### Planned Forage Crop and Seeding Rate (forage crop may be single species or mix of two or more species)

Species	Variety	Seed Size	Typic <mark>al</mark> Seeding D <mark>epth</mark>	Seeding Rate (PLS lbs/acre)	Percent of Mix (%)

## **Forage Crop Establishment and Management Considerations:**

- Establish forage crop mix as soon as practical prior to or after harvest of the production crop.
- During implementation, forage crop must not be grazed or burned.
- During implementation, notify NRCS of any planned changes in forage crop mix or crop rotation, or management to verify the planned system meets the enhancement criteria.

E328I - Forage harvest to reduce water	August 2019	
quality impacts by utilization of excess soil	Montana Supplement March 2020	Page   4
Nutrients with Montana Supplement		



• After implementation, if changes were made, update the tables above to document the applied crop rotation for the contract period and provide to NRCS.

### After implementation, complete the table below and provide to NRCS

Task	Provide information and details	
Seedbed Preparation		
Seeding Date		A
Seeding Depth		
Seeding Method		
Fertilizer, as needed		
Weed Management, as needed		
Harvest Date (window)		j
Harvest Method		

#### **NRCS will:**

- As needed, provide technical assistance in selecting forage crop for the crop rotation or substitute species that would meet the criteria of the enhancement. Forage crop may consist of a single species or mix of two or more species. Reference Montana Conservation Practice Standard, Specifications and Job Sheets for 328 and 340 in eFOTG.
- As needed, provide additional assistance to the participant as requested.
- Prior to implementation, verify the enhancement is being planned on fields where excess soil nutrients cause or increase water quality degradation concerns. Presence of excess nutrients must be identified in recent soil tests or increased risk to water quality documented by risk assessment tool. <REFER TO STATE SPECIFIC GUIDANCE>
- Prior to implementation, use information provided from the participant to calculate the
  average annual soil erosion value (water and wind) for each field using NRCS erosion
  prediction technologies.

Benchmark Management Soil Loss =	tons/acre/year
Planned Management Soil Loss =	tons/acre/year

• During implementation, evaluate any planned changes in forage crop selected, timing in crop rotation, management, or field operations to verify the new system meets the enhancement criteria.

E328I - Forage harvest to reduce water	August 2019	
quality impacts by utilization of excess soil	Montana Supplement March 2020	Page   5
Nutrients with Montana Supplement		



## **United States Department of Agriculture**

 After implementation, if there were any changes to planned rotation or management evaluate the applied crop rotation using information provided from the participant to calculate average annual erosion value to document that the applied rotation meets the enhancement criteria.

Applied Management Soil Loss = \_\_\_\_\_ tons/acre/year

#### **NRCS Documentation Review:**

I have reviewed all required participant documentation and have determined the participant has implemented the enhancement and met all criteria and requirements.

Participant Name	Contract Number
Total Amount Applied	Fiscal Year Completed
NRCS Technical Adequacy Signature	- Date

E328I - Forage harvest to reduce water quality impacts by utilization of excess soil Nutrients with Montana Supplement

August 2019

**Montana Supplement March 2020** 

Page | 6